

## Pretarsal Fixation of Gold Weights in Facial Nerve Palsy

Stuart R. Seiff, M.D., John H. Sullivan, M.D., L. Neal Freeman, M.D., and James Ahn, M.D.

Implantation of a gold weight in the pretarsal space was performed on 17 patients with poor eyelid closure secondary to facial nerve weakness. Successful results were obtained in 14 (82%) of 17 patients. Postmortem histopathology of such a case demonstrated firm fibrous encapsulation of the weight. Although some authors advocate fixation of lid weights to the orbital septum, fixation in the pretarsal space is preferred for the following reasons: (a) "cheesewiring" is less likely to occur, (b) a limited levator recession may be more easily performed, (c) greater mechanical benefits are obtained, and (d) the potential for inferior migration is less.

**Key Words:** Acoustic neuroma—Facial nerve palsy—Gold weight—Lagophthalmos—Lid loading—Seventh nerve palsy.

Function of the facial nerve may be lost on a temporary or permanent basis due to a variety of causes including tumor, trauma, infection, surgery, and idiopathic (including Bell's palsy). Ophthalmologists are often called on to manage problems such as lacrimal hyposecretion, brow ptosis, lagophthalmos, lid retraction, paralytic ectropion, and exposure keratopathy in these patients.


Various measures are available for treatment depending upon the nature and severity of the problem. Lagophthalmos may result in a medically unresponsive exposure keratopathy and require surgical correction. A lateral tarsal strip procedure may be done (1), but additional measures are usually necessary. Tarsorrhaphy is commonly performed; however, this limits visual field and may obscure the visual axis (2). The implantation of various different devices has been used in order to provide a more dynamic cure than that provided by tarsorrhaphy. Sheehan described the use of a tantalum wire and mesh implant in the upper lid (3). Fourteen years later, Morel-Fatio and Lalardrie reported on the use of a palpebral spring placed in the upper lid; this may be particularly useful in patients with severe impairment of eyelid closure (2,4). Use of an encircling silicone rod is another approach that may be beneficial in some patients (5). Techniques such as direct facial nerve repair, autogenous nerve grafting, cross-face nerve grafting, and nerve crossovers (especially utilizing the hypoglossal nerve) may be employed. Muscle transfers, free muscle grafts, and suspension with fascia lata, tendon, or alloplastic materials help provide bulk and support to the paralytic face (6).

The implantation of a gold weight to allow for gravity-assisted closure of the upper lid was first described by Smellie (7). Others have subsequently reported on their experience with gold weight implantation (2,8-14). Many of these authors believe the weight should be attached to the orbital septum

---

From the Ophthalmic Plastic and Reconstructive Surgery Service, Department of Ophthalmology, University of California, San Francisco.

Address correspondence and reprint requests to Stuart R. Seiff, M.D., Ophthalmic Plastic and Reconstructive Surgery Service, Department of Ophthalmology, A-750, University of California School of Medicine, San Francisco, CA 94143, U.S.A.



**FIG. 1.** Preoperative determination of appropriate weight. The correct weight places the upper lid 2 mm below the superior limbus with the eye open (a), and allows for passive closure (b).

(2,9,12,13). We describe our technique of implantation and attachment of the gold weight in the pretarsal space combined with limited levator recession, and report the results we have obtained in a series of patients. Histopathology is available for one case.

#### PATIENTS AND METHODS

The records of the surgical practices of two of us (S.R.S. and J.H.S.) were reviewed to identify patients who had undergone implantation of a gold weight. Seventeen patients were identified, each of whom required gold weight implantation because of ophthalmologic complications of facial nerve palsy.

The indications for surgery were essentially those outlined by Levine (15). Patients were strongly considered for gold weight loading if facial paralysis was likely to be longstanding and if keratopathy and/or marked irritation was present despite intensive medical measures or previous surgery. Surgery was more likely to be performed if decreased tear production, loss of corneal sensation, and/or poor Bell's phenomenon were present.

The gold weights are fabricated from 99.99% pure gold and are curved to fit the shape of the upper lid. The dimensions are approximately 10 mm length, 5 mm height, and 1 mm thickness. Two or three 1 mm holes are spaced across the length of the weight to allow for suture fixation. Weights are available in 0.2 g increments varying between 0.6 and 1.6 g.

The appropriate weight was selected for each patient preoperatively. By taping the different weights to the outside of the lid in the area overlying the tarsus, the correct weight for a given patient could be selected by a trial and error basis. The desired weight placed the upper lid 2 mm below the superior limbus with the eye open, and allowed the lid to close passively with the patient in the sitting position (Fig. 1). A slight overcorrection is desir-


able since the levator muscle appears to strengthen postoperatively.

The surgical technique is as follows: Local anesthesia is achieved with the use of topical 0.5% proparacaine hydrochloride and subcutaneous injection of 1.5 cc of 2% lidocaine with 1:100,000 epinephrine into the upper lid. The lid crease is marked. A 4-0 black silk suture is placed near the upper lid margin to allow for downward traction. A blade is used to incise at the lid crease through skin and orbicularis. Scissors dissection is performed inferiorly to the pretarsal space. The levator aponeurosis is stripped from its attachments to tarsus in the area of planned implantation, thus baring the anterior tarsal surface and effecting a modest levator recession. Hemostasis is achieved. The previously selected gold weight (which has been thoroughly cleansed and sterilized) is centered over the bare tarsal surface, and sutured directly to the tarsus with three interrupted 5-0 polyester sutures (Fig. 2). Antibiotic solution is irrigated into the



**FIG. 2.** The gold weight is centered over the bare tarsal surface and sutured directly to the tarsus.

FIG. 5. A dense layer of fibrous tissue (asterisk) surrounds the space where the gold implant had been previously placed (star). T, tarsus.



third patient had persistent corneal staining despite use of the lid load and eventually required additional tarsorrhaphy.

Special mention should be made of two patients who were considered successes. In one case, there was good closure postoperatively and the patient was comfortable, but ptosis was noted about 3 months following gold weight implantation. This was thought to be due to partial return of facial nerve function plus the effect of the lid load. The gold weight was removed and ptosis repair was subsequently carried out. Another patient initially underwent implantation of a 1.6 g weight; however, wound closure was modified from that described above. In this case, supratarsal fixation of the levator aponeurosis to the skin was included in an effort to form a defined lid crease. The gold weight began to extrude 2 months later and was removed. One month later, a 1.4 g weight was implanted with routine skin closure without supratarsal fixation. The patient has done well since the second operation. No other cases of extrusion occurred in this series.

Histopathology was obtained from an 81-year-old man who underwent gold lid loading for facial nerve palsy caused by *Pseudomonas* otitis. The postoperative result was good with excellent lid closure and a clear cornea with minimal or no staining. Four months after the lid surgery, the patient died from a myocardial infarction and permission for autopsy was granted.

Histopathologic examination revealed a dense layer of fibrous tissue surrounding the area in the

pretarsal space where the gold implant had been placed (Fig. 5). A few lymphocytes and epithelioid cells were seen in the region of the fibrous tissue. No foreign body giant cells were present.

## DISCUSSION

Tarsorrhaphy has been traditionally used when exposure keratopathy develops in patients with facial nerve palsy, but besides limiting vision, the procedure may lead to an unappealing cosmetic defect. Following release of a tarsorrhaphy, problems such as lid notching or entropion may occur (2).

Lid loading operations may be very effective in patients with weakness of the facial nerve. Gold has been used more commonly than other materials because of superior color match as seen through skin combined with minimal tissue reactivity (7). The surgical technique of lid loading is relatively simple, and by selecting the correct weight for loading, the eyelids will open and close well on volition. Though dependent upon gravity for its effect, closure in the supine position (e.g., during sleep) is possible if the head is slightly elevated (13). If facial nerve function returns or complications associated with the lid weight develop, the weight may be easily removed.

Different authors have described various locations in which the lid load may be implanted. Smellie indicated placement of the weight subcutaneously in the lower part of the lid without mentioning whether dissection was superficial or deep

5. Arion HG. Dynamic closure of the lids in paralysis of the orbicularis muscle. *Int Surg* 1972;57:48-50.
6. Baker DC. Facial paralysis. In: Smith BC, Della Rocca RC, Nesi FA, Lisman RD, eds. *Ophthalmic plastic and reconstructive surgery*. St. Louis: C. V. Mosby, 1987:580-90.
7. Smellie GD. Restoration of the blinking reflex in facial palsy by a simple lid-load operation. *Br J Plast Surg* 1966;19:279-83.
8. Barclay TL, Roberts AC. Restoration of movement to the upper eyelid in facial palsy. *Br J Plast Surg* 1969;22:257-61.
9. Jobe RP. A technique for lid loading in the management of the lagophthalmos of facial palsy. *Plast Reconstr Surg* 1974;53:29-32.
10. Habal HB. On lid loading in the management of lagophthalmos. *Plast Reconstr Surg* 1974;54:211.
11. Micheli-Pellegrini D. More on lid-loading in the management of lagophthalmos. *Plast Reconstr Surg* 1975;55:482.
12. Freeman BS. Facial palsy. In: Converse JM, ed. *Reconstructive plastic surgery*, 2nd ed. Philadelphia: W. B. Saunders, 1977:1835-8.
13. May M. *The facial nerve*. New York: Thieme, Inc., 1986:769-70.
14. Jackson IT. Surgical treatment of eyelid problems in facial palsy. In: Tessier P, Callahan A, Mustarde JC, Salyer KE, eds. *Symposium on plastic surgery in the orbital region*. St. Louis, C. V. Mosby, 1986:389-403.
15. Levine RE. Eyelid reanimation surgery. In: May M. *The facial nerve*. New York: Thieme, Inc., 1986:681-94.